

Drug Facts

Active ingredients

(in each 5 mL teaspoonful)

Aluminum hydroxide 200 mg (equivalent to dried gel USP).....	Antacid
Magnesium hydroxide 200 mg.....	Antacid
Simethicone 20 mg.....	Antigas

Purposes

Uses relieves • heartburn • sour stomach • acid indigestion
• the symptoms referred to as gas

Warnings

Ask a doctor before use if you have • kidney disease
• a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a
prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children.

Directions • shake well before use • do not take more than 24
teaspoonful in 24 hours or use the maximum dosage for more than 2
weeks • adults and children 12 years and older: take 2 to 4
teaspoonful between meals, at bedtime, or as directed by a doctor
• children under 12 years: Do not use

Other information • each 5 mL teaspoonful contains:
magnesium 85 mg, sodium 1 mg • store at room temperature
• protect from freezing • keep tightly closed • **TAMPER-EVIDENT:**
Do not use if breakaway band on bottle cap is missing or broken.

Inactive ingredients benzyl alcohol, butylparaben, carboxy-
methylcellulose sodium, flavor, hypromellose, microcrystalline cellulose,
propylparaben, purified water, saccharin sodium, sorbitol solution

*Major® Pharmaceuticals is not affiliated with the owner of the registered
trademark MYLANTA®.

Distributed by

MAJOR PHARMACEUTICALS

31778 Enterprise Drive,
Livonia, MI 48150 USA
TO REPORT ADVERSE
DRUG EVENTS CALL:
800-616-2471

Re-Order No. 002668

Rev 11/10 M-99 2-926-06112-5

