

B4423


Drug Facts
KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Uses for the prevention and treatment of nausea, vomiting, and dizziness associated with motion sickness

Active ingredient (in each tablet) Dimenhydrinate 50 mg Antiemetic

Warnings Do not use for children under 2 years of age unless directed by a doctor.


Drug Facts (continued)

 **MAJOR**[®]

NDC 0904-2051-12
Compare to the Active Ingredient in **DRAMAMINE**[®] Original Formula*

Driminate[™]
Dimenhydrinate USP, Antiemetic

For Nausea, Dizziness and Vomiting from Motion Sickness



**12 Tablets
50 mg EACH**
Use as Directed

* This product is not manufactured or distributed by Prestige Brands, Inc., owner of the registered trademark, Dramamine[®] Original Formula. 50844 REV1115K19802
Distributed by **MAJOR PHARMACEUTICALS**
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152 USA
Re-order No. 700466
M-17 Rev.09/16

TAMPER EVIDENT DO NOT USE IF PACKAGE IS OPENED OR IF PLESTER OF BLENDED CAPSULES SHOWS ANY SIGNS OF TAMPERING

Drug Facts (continued)

Inactive ingredients croscarmellose sodium, ■ see end flap for expiration date and lot number
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ protect from moisture

Drug Facts (continued)

Questions or comments? (800) 616-2471

cellulose, silica gel, stearic acid
dicalcium phosphate, magnesium stearate, microcrystalline

B-1212-198-02
REV1115K19802

Drug Facts (continued)

Ask a doctor before use if you have
■ a breathing problem such as emphysema or chronic bronchitis
■ glaucoma
■ difficulty in urination due to enlargement of the prostate gland or transurethral prostatectomy
■ Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product
■ avoid alcoholic beverages
■ avoid alcohol
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ use caution when driving a motor vehicle or operating machinery
■ If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Drug Facts (continued)

Directions
■ to prevent motion sickness, the first dose should be taken one-half to one hour before starting activity
■ adults and children 12 years and over: 1 to 2 tablets every 4-6 hours; do not exceed 8 tablets in 24 hours, or as directed by a doctor
■ children 6 to 12 years: 1½ to 2 tablets every 6-8 hours; do not exceed 3 tablets in 24 hours, or as directed by a doctor
■ children 2 to 6 years: ½ tablet every 6-8 hours; do not exceed 1½ tablets in 24 hours, or as directed by a doctor

Other information
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLENDED CAPSULE IS TORN OR BROKEN

3 09042 05112 5

No print/No varnish
Lot & Exp date